

## International Journal of Advanced Biochemistry Research



ISSN Print: 2617-4693  
 ISSN Online: 2617-4707  
 IJABR 2024; 8(6): 619-626  
[www.biochemjournal.com](http://www.biochemjournal.com)  
 Received: 04-05-2024  
 Accepted: 10-06-2024

**Paromita Mukherjee**  
 Researcher, Department of  
 Food and Nutrition, Swami  
 Vivekananda University,  
 Barrackpore, West Bengal,  
 India

**Souvik Tewari**  
 Assistant Professor,  
 Department of Food and  
 Nutrition, Swami Vivekananda  
 University, Barrackpore, West  
 Bengal, India

**Anindita Ray Chakravarti**  
 Assistant Professor and Head,  
 Department of Food and  
 Nutrition, Maharani Kasiswari  
 College, University of Calcutta,  
 Kolkata, West Bengal, India

**Corresponding Author:**  
**Souvik Tewari**  
 Assistant Professor,  
 Department of Food and  
 Nutrition, Swami Vivekananda  
 University, Barrackpore, West  
 Bengal, India

## Development of novel non-dairy synbiotic chocolates for promoting desirable gut microbiome

**Paromita Mukherjee, Souvik Tewari and Anindita Ray Chakravarti**

**DOI:** <https://doi.org/10.33545/26174693.2024.v8.i6h.1389>

### Abstract

This research focuses on the development and analysis (physico-chemical analysis and probiotic viability count) of non-dairy synbiotic chocolates, targeting consumers who are lactose intolerant, vegans, or seeking health-enhancing food options. The formulations were enriched with almond milk, oats powder, beetroot powder and *Streptococcus thermophilus* BURD PB 8 (Accession number: MN121741) to enhance both nutritional value and sensory properties. The proximate composition and antioxidant properties of the chocolates were evaluated. The results showed that T<sub>0</sub> had the highest carbohydrate content (42.88±0.01), significantly higher than T<sub>1</sub> (24.62±0.01) T<sub>2</sub> (20.38±0.01), and T<sub>3</sub> (15.65±0.01). Total fat content was substantially higher in T<sub>0</sub> (39.89±0.89) compared to T<sub>1</sub> (4.69±0.30), T<sub>2</sub> (4.42±0.26), and T<sub>3</sub> (3.76±0.33). The antioxidant activity, measured as DPPH radical scavenging activity (%), varied significantly among the treatments. At the beginning, the probiotic viability count of experimental chocolates was 3.80×10<sup>8</sup> CFU/g. After 60 days of storage, the count dropped to 3.20×10<sup>8</sup> CFU/g. On day 30, the number of cells had decreased to less than 1000. These findings suggest that the incorporation of *Streptococcus thermophilus* BURD PB 8 with almond milk, oats powder, beetroot powder and stevia powder enhance the nutritional profile and enhances antioxidant activities, making these chocolates a valuable addition to the functional food market.

**Keywords:** Almond milk, Beetroot powder, chocolate, non-dairy products, probiotics, *Streptococcus thermophilus*, synbiotics

### Introduction

Functional foods are gaining popularity due to their potential health benefits beyond basic nutrition (Hasler, 2022) [7]. Synbiotic products, which combine probiotics and prebiotics (Sekhon and Jairath, 2010) [10], are particularly valued for their role in promoting gut health (Maffei, 2019) [8]. Basically, probiotics are live microorganisms that are good for human health. By altering the microflora, secreting antibacterial compounds, competing with pathogens and harmful microorganisms to stop their adhesion, scavenging nutrients required for pathogen survival, creating an antitoxin effect, and partially reversing the effects of infection on the gut epithelium, probiotics have an antimicrobial effect (Tewari *et al.*, 2019) [12]. In contrast, prebiotics are indigestible substances that help the colon by promoting the growth and activity of one or a few species of bacteria that are already there (Tewari *et al.*, 2023) [13].

This research focuses on the development and analysis (Physico-chemical analysis and probiotic viability count) of non-dairy synbiotic chocolates that have both probiotics and prebiotics, targeting consumers who are lactose intolerant, vegan, or seeking health-enhancing food options.

### Materials and Methods

This study was conducted at the department of Food and Nutrition, Swami Vivekananda University, Barrackpore, W.B., India.

In this research Non-dairy Synbiotic Chocolates were prepared according to standard Chocolate-making procedures, with the functional ingredients incorporated during the mixing stage. All raw materials were procured from local commercial suppliers of Behala, Kolkata, W.B., India. The control Chocolate (T<sub>0</sub>) was prepared by using cocoa butter, cocoa liquor, whole milk powder, sugar and soy lecithin (Talbot, 2009) [11]. The quantities of different ingredients for the preparation of control Chocolate are tabulated in Table 1

(Talbot, 2009) [11]. In case of experimental Non-dairy Synbiotic Chocolate preparation, oats powder, almond milk, beetroots powder, stevia powder and probiotic microorganisms (*Streptococcus thermophilus* BURD PB 8) were used in different proportion. This new probiotic strain *Streptococcus thermophilus* BURD PB 8 (Accession number: MN121741) was isolated and identified from local

curd sample from Purba Bardhaman district of West Bengal, India (Bandyopadhyay *et al.*, 2019) [4].

The newly developed Non-dairy Synbiotic Chocolate formulations were tempered, molded and stored at 4 °C. The quantities of different ingredients for the preparation of experimental Chocolates are tabulated in Table 2.

**Table 1:** Treatment combination of Control Chocolate (T<sub>0</sub>).

Treatments (Control chocolate)	Cocoa powder (%)	Cocoa liquor (%)	Whole milk powder (%)	Sugar Powder (%)	Soy lecithin (gm)
T <sub>0</sub>	20	12.4	25.4	43	0.5

In order to prepare the control kulfi (T<sub>0</sub>), various percentages of raw components were mixed together to ensure equal blending, including cocoa butter, cocoa liquor, whole milk powder, sugar, and soy lecithin. Following the

blending process, the kulfi was moulded after tempering at 45 °C. Cooling was then carried out at between 10 and 15 degrees Celsius (Talbot, 2009) [11].

**Table 2:** Treatment combination of newly developed Non-dairy Synbiotic Chocolates (T<sub>1</sub>, T<sub>2</sub> and T<sub>3</sub>)

Treatments (Experimental Chocolate)	Almond milk (mL)	Oats Powder (gm)	Beetroots Powder (gm)	Probiotics ( <i>Streptococcus thermophilus</i> BURD PB 8) in percentage	Coconut Oil (mL)	Stevia Powder (gm)	Soy lecithin (gm)
T <sub>1</sub>	60	30	7	1	2	0.10	0.5
T <sub>2</sub>	65	25	5	1	4	0.50	0.5
T <sub>3</sub>	70	20	3	1	6	1	0.5

The experimental chocolate formulations detailed in the table vary in their composition of key ingredients to enhance both nutritional value and sensory properties. Treatment T<sub>1</sub> includes 60 mL of almond milk, 30 g of oats powder, 7 g of beetroot powder, 1% probiotics, 2 mL of coconut oil, 0.10 g of stevia powder, and 0.5 g of soy lecithin. Treatment T<sub>2</sub> increases the almond milk to 65 mL and coconut oil to 4 mL, while reducing oats and beetroot powders to 25 g and 5 g respectively, with stevia powder increased to 0.50 g, maintaining the same probiotic and soy lecithin levels. Treatment T<sub>3</sub> further adjusts these proportions with 70 mL of almond milk, 20 g of oats powder, 3 g of beetroot powder, 6 mL of coconut oil, and 1 g of stevia powder, keeping the probiotics and soy lecithin constant. These variations are designed to explore the optimal balance for a healthful, palatable chocolate product.

### Analysis of product

Moisture Content (%), Total Ash (%), Protein Content (%), Total Fat (%), Dietary Fibre (%), p<sup>H</sup>, Total soluble Solid (Brix) and DPPH (%) were estimated by AOAC (2000) [12]

method. And Carbohydrate Content (%) was calculated by difference. To find out the Probiotic Viability count (CFU/g) of the final product, suitable dilutions were poured with respective agar media in sterile Petri dishes in duplicates. In addition to viable count of *Streptococcus thermophilus* BURD PB 8 on MRS agar (after incubation at 37 °C/48 h under facultative anaerobic conditions) (Auty *et al.*, 2001) [3].


### Results

This study evaluates the proximate composition and antioxidant content of different treatments (T<sub>0</sub>, T<sub>1</sub>, T<sub>2</sub>, and T<sub>3</sub>) of a food product. The analyzed parameters include moisture, total ash, carbohydrates, protein, total fat, dietary fibre, p<sup>H</sup>, Total soluble Solid and antioxidant percentages. Each parameter was assessed for significant differences across treatments. The values are presented as means ± standard deviations. Significant differences between treatments are indicated by different superscript letters (abcd).

**Table 3:** Physicochemical analysis of control and experimental non-dairy synbiotic chocolates

Parameters	T <sub>0</sub>	T <sub>1</sub>	T <sub>2</sub>	T <sub>3</sub>
Moisture (%)	2.60±0.01 <sup>abcd</sup>	2.17±0.01 <sup>bd</sup>	2.21±0.01 <sup>c</sup>	2.24±0.01 <sup>d</sup>
Total ash (%)	4.06±0.01 <sup>abcd</sup>	6.47±0.11 <sup>bcd</sup>	5.94±0.01 <sup>c</sup>	4.96±0.01 <sup>d</sup>
Carbohydrate (%)	42.88±0.01 <sup>abcd</sup>	24.62±0.01 <sup>bd</sup>	20.38±0.01 <sup>cd</sup>	15.65±0.01 <sup>d</sup>
Protein (%)	4.06±0.017 <sup>abc</sup>	6.47±0.01 <sup>bd</sup>	5.94±0.012 <sup>c</sup>	4.96±0.01 <sup>d</sup>
Total Fat (%)	39.89±0.89 <sup>abcd</sup>	4.69±0.30 <sup>b</sup>	4.42±0.26 <sup>c</sup>	3.76±0.33 <sup>d</sup>
Dietary Fibre (%)	1.85±0.02 <sup>abcd</sup>	3.26±0.03 <sup>bcd</sup>	2.71±0.04 <sup>cd</sup>	2.18±0.04 <sup>d</sup>
p <sup>H</sup>	6.10±0.05 <sup>acd</sup>	6.30±0.05 <sup>b</sup>	6.40±0.05 <sup>c</sup>	6.50±0.05 <sup>d</sup>
Total Soluble Solid (Brix)	44.70±0.11 <sup>abcd</sup>	42.80±0.05 <sup>bcd</sup>	42.40±0.11 <sup>cd</sup>	42.00±0.11 <sup>d</sup>
DPPH (%)	52.0±0.05 <sup>abcd</sup>	72.5±0.02 <sup>bcd</sup>	68.3±0.05 <sup>cd</sup>	64.1±0.05 <sup>d</sup>


All the test were performed in triplets. Different letter in the same column indicates statistical significance level of p<0.0001.



**Fig 1:** Graphical representation of moisture content (%) of control and experimental non-dairy synbiotic chocolates

The moisture content varied among the treatments, with T<sub>0</sub> (2.600±0.01) having the highest moisture content, followed by T<sub>3</sub> (2.240±0.01), T<sub>2</sub> (2.210±0.01), and T<sub>1</sub> (2.170±0.01). The differences were statistically significant, as indicated by


the superscript letters. Lower moisture content in T<sub>1</sub>, T<sub>2</sub>, and T<sub>3</sub> compared to T<sub>0</sub> suggests that these treatments are more stable and less prone to microbial spoilage, enhancing shelf life.



**Fig 2:** Graphical representation of total ash (%) of control and experimental non-dairy synbiotic chocolates

Total ash content, an indicator of mineral content, was highest in T<sub>1</sub> (6.470±0.11), followed by T<sub>2</sub> (5.940±0.01), T<sub>3</sub> (4.960±0.01), and T<sub>0</sub> (4.060±0.01). The significant increase in ash content from T<sub>0</sub> to T<sub>1</sub> and T<sub>2</sub> suggests that these


treatments have a higher mineral content, which could contribute to improved nutritional value. However, the slightly lower ash content in T<sub>3</sub> indicates a decrease in mineral content compared to T<sub>1</sub> and T<sub>2</sub>.



**Fig 3:** Graphical representation of carbohydrate (%) of control and experimental non-dairy synbiotic chocolates

Carbohydrate content showed a decreasing trend from T<sub>0</sub> to T<sub>3</sub>. T<sub>0</sub> had the highest carbohydrate content (42.88±0.017), significantly higher than T<sub>1</sub> (24.62±0.011), T<sub>2</sub> (20.38±0.011), and T<sub>3</sub> (15.65±0.017). This reduction in


carbohydrate content across treatments suggests a formulation change that replaces carbohydrates with other components, potentially aiming to create a lower-carb product suitable for specific dietary needs.



**Fig 4:** Graphical representation of protein content (%) of control and experimental non-dairy synbiotic chocolates

Protein content was highest in T<sub>1</sub> (6.470±0.011) and T<sub>2</sub> (5.940±0.012), significantly higher than T<sub>0</sub> (4.060±0.017) and T<sub>3</sub> (4.960±0.011). The increased protein content in T<sub>1</sub> and T<sub>2</sub> suggests these treatments were designed to enhance


the protein profile of the product, making them more suitable for consumers seeking higher protein intake, such as athletes and bodybuilders.



**Fig 5:** Graphical representation of total fat (%) of control and experimental non-dairy synbiotic chocolates

Total fat content was substantially higher in T<sub>0</sub> (39.89±0.89) compared to T<sub>1</sub> (4.693±0.30), T<sub>2</sub> (4.423±0.26), and T<sub>3</sub> (3.763±0.33). The significant reduction in fat content from


T<sub>0</sub> to the other treatments indicates an effort to prepare a lower-fat product. This makes T<sub>1</sub>, T<sub>2</sub>, and T<sub>3</sub> more suitable for health-conscious consumers aiming to reduce fat intake.



**Fig 6:** Graphical representation of dietary fibre (%) of control and experimental non-dairy synbiotic chocolates

Dietary fibre content was highest in T<sub>1</sub> (3.260±0.03) followed by T<sub>2</sub> (2.717±0.04) and T<sub>3</sub> (2.183±0.04), with T<sub>0</sub> having the lowest fibre content (1.850±0.02). The increase in dietary fibre in T<sub>1</sub>, T<sub>2</sub>, and T<sub>3</sub> indicates these formulations


are designed to enhance fibre intake, which is beneficial for digestive health and can help in managing weight and blood sugar levels.



**Fig 7:** Graphical representation of pH of control and experimental non-dairy synbiotic chocolates

The pH values of the Non-Dairy Synbiotic Chocolates varied significantly among the different treatments (T<sub>0</sub>, T<sub>1</sub>, T<sub>2</sub>, T<sub>3</sub>). The control sample (T<sub>0</sub>) exhibited the lowest pH of 6.1±0.05, while T<sub>3</sub> had the highest pH of 6.5±0.05. The gradual increase in pH from T<sub>0</sub> to T<sub>3</sub> suggests that the addition of almond milk, oats powder, and *Streptococcus thermophilus*


BURD PB 8 influenced the acidity of the chocolate. Specifically, the fermentation process likely contributed to this pH increase as the probiotic bacteria metabolized substrates, potentially producing less acidic by products compared to the original mixture.



**Fig 8:** Graphical representation of Total Soluble Solid (Brix) of control and experimental non-dairy synbiotic chocolates

The total soluble solids, measured in degrees Brix (°Brix), also showed significant differences across the treatments. The control sample (T<sub>0</sub>) had the highest °Brix value at 44.7 ± 0.11, while the synbiotic chocolate with the highest probiotic content (T<sub>3</sub>) had the lowest °Brix value at 42.0±0.11. This decrease in °Brix from T<sub>0</sub> to T<sub>3</sub> indicates

that the fermentation by *Streptococcus thermophilus* BURD PB 8 consumed some of the soluble solids, likely sugars, resulting in a lower °Brix value. This aligns with the expected metabolic activity of probiotics, which utilize available sugars for growth and activity.



**Fig 9:** Graphical representation of DPPH (%) of control and experimental non-dairy synbiotic chocolates

The antioxidant activity, measured as DPPH radical scavenging activity (%), varied significantly among the treatments. The control sample (T<sub>0</sub>) had the lowest DPPH activity at 52.0±0.05%, whereas T<sub>1</sub> exhibited the highest activity at 72.5±0.02%. The antioxidant activity decreased slightly in T<sub>2</sub> (68.3±0.05%) and further in T<sub>3</sub> (64.1±0.05%). The increase in antioxidant activity from T<sub>0</sub> to T<sub>1</sub> suggests that the initial addition of almond milk, oats powder, and the probiotic improved the antioxidant properties of the chocolate. A similar outcome from a previously published study showed that the use of Galacto-oligosaccharides (GOS) increased the DPPH (%) in chocolate (Tewari *et al.*, 2023) [13]. However, the slight decrease in DPPH activity in T<sub>2</sub> and T<sub>3</sub> could indicate that beyond a certain concentration, the components might not synergize as effectively to boost antioxidant properties, or that the fermentation process slightly reduces certain antioxidant compounds.

#### Probiotic Viability:

The initial CFU count was 3.80×10<sup>8</sup> CFU/g. The count declined to 3.20×10<sup>8</sup> CFU/g during the first 60 days of storage. However, the cell count had fallen below 1000 on the 30th day. To confirm this observation, 1 g of chocolate was suspended in 100 ml sterile skim milk and was incubated at 37 °C. After 24 h, the milk was found in coagulated form having 0.8% lactic acid. The cells were not capable to form colonies while plated on MRS. It is difficult to give the exact reason for this, but it could be attributed to the complex internal structure of chocolate. The findings of this study are close to those of Gadhiya *et al.* (2018) [6].

#### Discussion

The developed non-dairy synbiotic chocolates have advantageous physicochemical characteristics, such as a high protein content that aids in muscle maintenance and repair and a low moisture level that extends shelf life. The

fat content supplies important fatty acids and is mostly sourced from almond milk (Vanga and Raghavan, 2018) [15]. The use of oats powder significantly increases dietary fibre content, promoting digestive health (Tosh and Bordenave, 2020) [14].

The observed increase in p<sup>H</sup> values with the addition of *Streptococcus thermophilus* BURD PB 8 can be attributed to the metabolic activities of the probiotic, which may produce less acidic byproducts during fermentation. This rise in p<sup>H</sup> indicates a shift in the chemical environment of the chocolate, which could affect its flavor and texture (Afoakwa *et al.*, 2008) [1].

The decrease in total soluble solids (°Brix) across the treatments demonstrates the effective fermentation by the probiotic, which utilizes soluble components for growth. This is a desirable outcome in synbiotic products as it signifies active fermentation, contributing to the health benefits associated with probiotics (Markowiak and Śliżewska, 2017) [9].

The significant enhancement in DPPH radical scavenging activity in T<sub>1</sub> compared to T<sub>0</sub> highlights the beneficial impact of adding almond milk, oats powder, and probiotics on the antioxidant properties of the chocolate. The slight reduction in antioxidant activity in T<sub>2</sub> and T<sub>3</sub> suggests a potential threshold effect, where optimal concentrations of added components maximize antioxidant properties, beyond which the benefits plateau or decrease.

Overall, the results indicate that the incorporation of almond milk, oats powder, and *Streptococcus thermophilus* BURD PB 8 in non-dairy synbiotic chocolates enhances their functional properties, particularly in terms of p<sup>H</sup> stability, reduction in soluble sugars, and improved antioxidant activity, making them a potentially healthier alternative to traditional dairy chocolates.

These findings suggest that the synbiotic chocolates can serve as a nutritious, functional food suitable for various



consumer groups, including those with lactose intolerance, vegans, and health-conscious individuals. The incorporation of almond milk, oats powder, and *Streptococcus thermophilus* BURD PB 8 not only improves the nutritional profile but also offers additional health benefits, making these chocolates a valuable addition to the functional food market.

### Conclusion

The physicochemical analysis of the newly developed non-dairy synbiotic chocolates enriched with almond milk, oats powder, and *Streptococcus thermophilus* BURD PB 8 demonstrates their potential as a functional food product. The enhanced nutritional profile and probiotic benefits make these chocolates suitable for health-conscious consumers and those with specific dietary requirements.

### Acknowledgments

All the authors would like to express their sincere thanks to the Department of Food and Nutrition, Swami Vivekananda University (Barrackpore, West Bengal, India) and Oriental Institute of Science and Technology Vidyasagar University (Dewandighi, Burdwan, West Bengal, India) for providing research facilities. Authors also grateful to Dr. Bidyut Bandyopadhyay and Atrayee Roy in achieving the research objective.

**Conflicts of interest:** The authors declare no conflict of interest.

### References

1. Afoakwa EO, Paterson A, Fowler M, Ryan A. Flavor formation and character in cocoa and chocolate: a critical review. *Crit Rev Food Sci Nutr*. 2008;48(9):840-857.
2. AOAC International. Official Methods of Analysis of AOAC International. 17<sup>th</sup> ed. Gaithersburg, MD: AOAC International; c2000. ISBN: 978-0935584677.
3. Auty MAE, Gardiner GE, McBrearty SJ, O'Sullivan EO, Mulvihill DM, Collins JK, *et al*. Direct in situ viability assessment of bacteria in probiotic dairy products using viability staining in conjunction with confocal scanning laser microscopy. *Appl Environ Microbiol*. 2001;67(1):420-425.
4. Bandyopadhyay B, Roy A, Maitra M. Phenotypic characterization and genotypic identification of local milk fermented *probiotics*. *Biotechnol Ind J*. 2019;15(4):194.
5. Barros L, Baptista P, Ferreira IC. Effect of *Lactarius piperatus* fruiting body maturity stage on antioxidant activity measured by several biochemical assays. *Food Chem Toxicol*. 2007;45(9):1731-1737.
6. Gadhiya D, Shah NP, Patel AR, Prajapati JB. Preparation and shelf life study of probiotic chocolate manufactured using *Lactobacillus helveticus* MTCC 5463. *Acta Aliment*. 2018;47(3):350-358.
7. Hasler CM. Functional foods: benefits, concerns and challenges—a position paper from the American Council on Science and Health. *J Nutr*. 2002;132(12):3772-3781.
8. Maftei NM. Probiotic, prebiotic and synbiotic products in human health. In: Talbot G, ed. Science and technology of enrobed and filled chocolate, confectionery and bakery products. Intech Open; c2019.

9. Markowiak P, Ślizewska K. Effects of probiotics, prebiotics, and synbiotics on human health. *Nutrients*. 2017;9(9):1021.
10. Sekhon BS, Jairath S. Prebiotics, probiotics and synbiotics: An overview. *J Pharm Educ Res*. 2010, 1(2).
11. Talbot G, ed. Science and technology of enrobed and filled chocolate, confectionery and bakery products. 2009.
12. Tewari S, David J, Gautam A. A review on probiotic dairy products and digestive health. *J Pharmacogn Phytochem*. 2019;8(3):368-372.
13. Tewari S, Pattanayak A, Zanwar S, Vaishnav S. Physico-chemical analysis of newly prepared prebiotic chocolates by using Galacto Oligosaccharides (GOS). *J Survey Fish Sci*. 2023;10(4S):1752-1763.
14. Tosh SM, Bordenave N. Emerging science on benefits of whole grain oat and barley and their soluble dietary fibers for heart health, glycemic response, and gut microbiota. *Nutr Rev*. 2020;78(Suppl 1):13-20.
15. Vanga SK, Raghavan V. How well do plant based alternatives fare nutritionally compared to cow's milk? *J Food Sci Technol*. 2018;55(1):10-20.